



Temporal and angular biomechanical analysis of the ankle joint as a basis for the prevention of external ligament tear injury in soccer players

*Layth Muthanna Ibrahim Al-Baitar*¹
*MOHAMED ABDULSALAM FADHIL*²
*MOHAMMED FAISAL MAHDI*³
layth.mu20@uosamarra.edu.iq
mohammed.abd.alsalam@uosamarra.edu.iq
mohammed.faisal@uosamarra.edu.iq

ABSTRACT

Objective: To analyse the temporal biomechanical characteristics and kinetic angles of the ankle joint during various high-risk situations in football (jumping and landing, cutting action, contact) to identify quantitative indicators associated with external ligament rupture injury.

The researcher used the descriptive-analytical method according to an intentional sample of (25) players from Samarra Club, aged 18-30 years and representing all positions. The performance was recorded with the help of a high-resolution digital camera and motion analysis employing Kinovea software in order to determine temporal variables (contact time, stability time) and angular variables (lateral deflection angle, angular velocity).

The results showed that:

- The time spent on the ground is relatively short (0.42 seconds) due to shock absorption efficiencies of most players.
- Allowed deflection angle varied (12°–27°), which implies uneven control of the joint. The angular velocity was high during the change of direction (245°/s), which increased tension on the ligaments.
- Post performed stabilization time (0.92-2.05 seconds), proves the ability of players to re-establish balance

Thus, thus the temporal and angular biomechanical analysis results are an accurate tool for evaluating players according to their risk levels (low-medium-high) and can establish the basis for implementing preventive programs based on dynamic balance exercises, strengthening of peroneal muscles and neuromuscular control trainings.

1-1 Introduction and Importance of the Research:

Ankle injuries are one of the most common injuries among soccer players, as studies indicate that ankle sprains and external ligament tears account for approximately 20% of sports injuries in high-level competitions, and often occur during jumping, landing, or sudden change of direction and physical contact, leading to prolonged absences of players and affecting athletic performance. (Buckthorpe et al., 2025, p. 214)

The problem with ankle injuries is not only their occurrence, but also their high frequency of occurrence, as a player who has suffered a previous injury becomes more likely to be injured again if it is not dealt with properly preventively, which illustrates the need for preventive programs based on a rigorous scientific analysis of movement. (Flore et al., 2024, pp. 33–34)

Temporal and angular biomechanical analysis of motion is an objective tool for understanding the mechanics of the ankle joint during high-risk situations such as jumping, landing, or changing direction. Through this analysis, quantitative variables such as contact time with the ground, and angular variables such as lateral deflection angle and rotational velocity, can be identified, which are key indicators for assessing the risk of injury.

(The Football Physio, 2024, p. 12)

Neuromuscular training programs such as FIFA 11+ have been shown to be effective in reducing ankle injuries among soccer players, improving dynamic balance, strengthening the muscles around the ankle, and enhancing neuromuscular control. However, the design of these programs must be based on accurate movement analytical data to ensure their effectiveness. (Eser et al., 2024, pp. 45–46) Dynamic balance training has also proven to play an important role in the prevention of ankle injuries, as it helps to improve functional stability and reduce neuromuscular response time during sudden situations in matches.

(Tomchuk et al., 2025, p. 88)

1-2 Research Problem:

Ankle injuries, especially external ligament tears, are one of the most common injuries in football, with UEFA reporting that these injuries account for 15-20% of all injuries recorded in major tournaments, and they often occur during jumps and landings, sudden change of direction or physical contact. (Ekstrand et al., 2019, p. 112)

Although there are many treatment and preventive programs, most of them rely on general protocols that are not based on an accurate biomechanical analysis of the actual movement of the players. This leaves a clear scientific gap in understanding the relationship between kinematic variables (e.g., contact time with the ground, angle of lateral deflection, angular velocity) and the probability of ankle injury.

(Delahunt et al., 2010, pp. 45–46)

Studies also show that players who have had a previous ankle injury are up to 70% more likely to re-injury within one year if preventive programs based on scientific analysis of movement are not applied, turning the injury into a chronic condition that affects a player's athletic career.

Hence, the research problem is identified in the absence of preventive programs based on temporal and angular biomechanical analysis of the ankle joint in high-risk situations in soccer players, which increases the likelihood of injury and recurrence of external ligament tears.

1-3 Research Objectives:

- Identify high-risk situations associated with the rupture of the external ligaments of the ankle joint in soccer players.
- Identify motor performance analysis using a single camera and free software (Kinovea) to monitor the temporal and angular variables of the ankle joint during these situations.

1 Practical implementation within this research.

1-4 Research Hypotheses:

1. There are statistically significant differences in the time of contact with the ground between high-risk situations (jumping and landing, changing direction, physical contact).
2. There are statistically significant differences in the angle of lateral deflection of the ankle joint between players with a history of ankle injury and those who have not suffered a previous injury.

1.5 Identification of Areas:

1.5.1 Human field: advanced Samarra FC players

1-5-2 Temporal Domain: For the period from (22/4/2025) to (2/8/2025).

1.5.3 Spatial Area: Samarra Sports Club Stadium.

2- Research methodology and field procedures:

2.1 Research Methodology:

The researcher used the descriptive-analytical method, supported by a controlled field experiment, by imaging the motor performance of the players during high-risk situations (jumping and landing, changing direction, physical friction), and analyzing the temporal and angular variables of the ankle joint using a single camera and free software, with the aim of extracting accurate quantitative indicators on which future preventive programs are based.

2.2 Research Population and Sample:

2.2.1 Research Community

The research community consists of footballers registered in the Iraqi Premier League and the First Division, as this community is the most susceptible to ankle injuries as a result of the intensity of matches and the frequency of high-risk situations.

2.2.2 Research Sample

An intentional sample was selected from 25 players from Samarra FC representing different positions (defenders, midfielders, forwards, goalkeepers), aged between 18 and

30 years, all of whom are in good physical fitness and have been free of a severe ankle injury during the last 6 weeks.

- Listing criteria: An officially registered player, capable of performing demo missions.
- Exclusion criteria: chronic ankle or knee injuries, use of therapeutic tools that alter the mechanics of movement.

The research population was selected from the second stage student at the Faculty of Physical Education and Sport Sciences, Samarra University, three study divisions for the academic year 2024-2025, which are (99), as the researcher selected the research sample consisting of (85) for the research necessities, and the failing and absent students were excluded for the homogeneity of the sample, as the sample was divided into (25) division for each of the three groups (individual experimental), (competitive experimental), (traditional control) and () students for the exploratory experiment, and Table (2) shows the division of the sample.

2.2.2.1 Sample homogeneity

In order to verify the suitability of the sample for biomechanical analysis, the researcher calculated the statistical homogeneity indices of the basic variables (age, height, weight, training age). The results showed that the mean and standard deviations came within the normal limits, and the coefficients of torsion and flattening were less than ± 1 , which indicates that the distribution of the data is close to the normal distribution. This confirms that the sample is homogeneous in terms of physical characteristics and training experience, so the differences that will emerge later can be attributed to experimental variables rather than to significant individual differences between players.

Table (1)
Sample homogeneity

N = 25

| Flattening coefficient | Torsion coefficient | Standard deviation | Arithmetic mean | Unit of Measurement | Variables |
|------------------------|---------------------|--------------------|-----------------|---------------------|--------------|
| -0.41 | 0.35 | 6.1 | 72.5 | kg | Mass |
| 0.27 | 0.27 | 5.2 | 176.8 | Poison | Length |
| 0.31 | 0.31 | 2.4 | 22.6 | Year | Age |
| -0.36 | 0.29 | 2.1 | 6.8 | Year | Training Age |

Table Interpretation:

- All statistical values are within the normal limits.
- The sample is homogeneous in terms of age, height, weight, and training age.
- This homogeneity enhances the credibility of the results and confirms that biomechanical analysis will reflect real differences in performance and not individual differences in basic characteristics.

2.2.2.2 Homogeneity of the sample in terms of injury

The researcher collected data on the history of ankle injuries among the sample members (25 players from Samarra Club), according to different positions (defenders, midfielders, forwards, goalkeepers) and classified them into three main categories:

- Players who have not suffered a previous ankle injury.
- Players who have suffered one ankle injury during their career.
- Players who have suffered repeated ankle injuries (more than twice).

Table (2)
Sample distribution by ankle injury date

N = 25

| Recurrent injuries (≥ 2) | One Injury | No injury | Number of Players | Center |
|---------------------------------|------------|-----------|-------------------|------------|
| 2 | 2 | 3 | 7 | Defender |
| 1 | 4 | 4 | 9 | Medium |
| 2 | 2 | 2 | 6 | Attacker |
| 1 | 1 | 1 | 3 | Goalkeeper |
| 6 | 9 | 10 | 25 | Total |
| 24% | 36% | 40% | 100% | Percentage |

Data provide in the table appears balanced among the centers, other than differences observed in rates of ankle injury. 40% were uninjured, 36% had had one injury and 24% recurrent injuries. In this case, defenders and attackers have an increased chance for a repetitive injuries owing to their continuous leverage with the ball and landing on opponent direction twisting towards it seems inevitable considering physical exertion levels cause more micro-trauma on muscle fiber that leads to cramping. Midfielders experience lesser number of injuries per match followed by goalkeepers whose count remains lower than all but are prone based on the type of objectives required from him (like frequent jumps etc.) leading them as significant members when assessing injury probability through team characteristics over time.

Such homogeneity in the distribution of injuries increases this credibility because it enables investigators to analyze the results on temporal and angular biomechanical variables and injury risk in players from different categories while ensuring that differences found are either real differences between motor performance or individual differences regarding history of disease.

2.2.2.3 Justification for sample selection

The sample of the research was chosen from the players of Samarra Club (25 players) for several scientific and applied considerations, the most important ones are as follows:

Ease of access and collaboration: The club is furnished with a one-stage training environment setting, administrative and technical collaboration thereby promoting orderly data collection and transfer of experience.

Homogenization of Physical and Training Characteristics: The homogeneity results are making result easy to interpret, because all players have similar age,height, weight, training age that is increasing power of biomechanical analysis.

Variety of positions : Players from different positions (defenders, midfielders, forwards, goalkeepers) are included in the sample enabling the study of injuries across various play settings.

Diverse injury history: Some players had never had an injury, while others had one or recurrent injuries, allowing for a comparison of biomechanical performance.

Relevancy to the local environment: Samarra Club was selected because it represents a microcosm of the industry reality in terms of resourced Iraqi clubs, such that outputs from this search will be directly applicable to the local context and potentially generalizable across similar clubs.

2.3 Tools and devices used for data collection:

2.3.1 Research tools

Foreign sources.

International Information Network.

2.3.2 Data collection methods

Kinovea Software.

A personal data form for writing down age, height, weight, training age, position played, dominant foot and history of ankle injury.

You are the performance and a of false seeking including to tables to release the values extracted from by degree (contact time, angle, angular velocity, stability time).

2.3.3 Field Research Tools

1. **4 Football Balls**
2. **2 cameras at 120 frames/second.**
3. **2 camera holders.**
4. **Tape measure.**

2.4 Defining Tests and Measurements

2.4.1 Defining the Search Test

The tests you used in your research were based on the typical scenarios related to ankle injury as it relates to football activity: jumping and landing, changing direction and physical contact with the ball. Jumping is one of the most common circumstances that cause ankle sprains, especially due to deficits in motor control or muscle fatigue as shown above from studies.

(Wright, Arnold, & Ross, 2016)

Video studies of professional football ankle injuries have similarly identified a cutting maneuver as a major risk factor for injury (5). (Kristianslund, Bahr, & Krosshaug, 2011) As stated by Owoeye, VanderWey, and Pike (2014), one of the most common ankle injuries occurring during match play is caused collateral impact for controlling a ball.

These tests were specifically chosen for clear scientific and practical reasons, rather than at random. They are realistic high-risk game scenarios, and the extraction of stroke details such as contact time, angle of deflection, angular velocity and stability time gives higher integrity for biomechanical analysis. Moreover, these tests can be utilized in the local environment with simple tools (a ball and a digital camera), while conserving Iraqi conditions suspected of being poor while maintaining scientific accuracy. (Faude, Rößler, & Junge, 2013)

In doing so, these tests provide a solid scientific basis to the analysis of motor performance in relation to injury risk and ensure that the responses obtained will be directly relevant in terms of training and pre-habilitation.

2.4.2 Identification of Biomechanical Variables

The biomechanical variables allowed the researchers to establish a relationship between these variables and the risk of ankle injury to investigate. These variables are: ground contact time, lateral deflection angle of the ankle, angular velocity in change of direction

and stability time after execution. These variables were also chosen because they are true objective measures that can be quantified with video analysis software like Kinovea, and have a history of use in biomechanical investigations to discriminate between dynamic stability (Wright, Arnold, & Ross, 2016).

Purpose: Contact time with the ground is a fundamental measure of shock absorption and motor control when landing; longer contact times are purported to increase injury risk (Bahr & Holme, 2003).

The ankle angle of lateral deflection represents the stability of the joint throughout performance, and video analyses demonstrate that increased deviations contribute to an increased rate of ankle sprains. (Kristianslund, Bahr, & Krosshaug, 2011)

The angular velocity of the ankle is crucial in change of direction to be completed with optimal control or minimizing load/ stress on joint ligaments as high speed is known to cause loading (Owoeye et al. 2014) Finally, the stabilization time following a test represents an indicator of the capacity of the neuromuscular system to recover balance, which is crucial in preventing recurrent injuries. (Faude, Rößler, & Junge, 2013)

Selecting these variables therefore not only captures generic measurements of anatomical structure and mechanical functioning but also mimics real-world play situations, supports a quantitative description of movement patterns that may be associated with ankle injury risk thereby improving the robustness/acceptance of results and rendering them applicable in training/prevention.

2.5 Main Experience:

The main experiment is designed to include three basic tasks that simulate the most common situations associated with ankle injuries in football: jumping and landing, changing direction, and physical contact with the ball. These tasks were carried out in a standardized training environment using a single ball and a high-resolution digital camera, with the aim of recording and biomechanically analyzing the motor performance.

In the first mission, the player jumps to pick up the ball with the head or control it with the chest, then lands on one foot while trying to maintain balance, a test that simulates playing situations that are associated with an increased risk of ankle sprains during a landing. The second task involves the player launching the ball at a medium speed and then changing direction at an angle of 45° or 90° while still controlling the ball, reflecting dribbling and abrupt cut situations that require high control of the ankle joint. In the third task, the player faces an opponent in a shoulder-to-shoulder position while trying to control or pass the ball, a test that simulates competitive situations in which the player is exposed to unexpected external pressure that increases the likelihood of injury.

This experiment was carried out according to a structured field protocol that ensures the standardization of conditions for all sample members. The pitch has been set up and the performance area has been defined so that the floor is level and secure, with the digital camera installed in a suitable side angle for full-action filming. The players also underwent a standard ten-minute warm-up program that includes light running and dynamic stretching exercises to reduce the likelihood of injuries during the experiment.

Each task was repeated three times to ensure stability, with uniform instructions for all players. All attempts were recorded using a digital camera, and then the videos were analyzed via Kinovea software to extract time and angle variables such as contact time, angle of deflection, angular velocity, and stability time. Finally, the extracted values were collected in special tables, reviewed for accuracy and error-free, in preparation for entering into the statistical program and later analyzing.

2.6 Statistical Method and Statistical Methods:

The research relied on the descriptive and analytical statistical method, with the aim of providing an accurate picture of the characteristics of the sample and the studied variables, and revealing the differences and biomechanical relationships associated with the risk of ankle injury.

In the descriptive aspect, a set of statistical methods such as arithmetic mean, standard deviation, torsion coefficient, and flattening coefficient were used to clarify the general characteristics of the sample and its distribution. In the analytical aspect, the t-test was applied to repeated measurements in order to compare the performance of the players in the three main tasks, in addition to using ANOVA) to detect the differences between the different positions within the team. The Pearson correlation coefficient was also used to study the relationships between biomechanical variables, such as the relationship between angular velocity and stability time, or between angle of deflection and contact time.

In order to provide a clearer picture of the risk levels, the internal classification method was adopted to divide the players into risk layers (low-medium-high) according to the extracted values, allowing the results to be used in building future prevention and training programs.

3. Presentation and discussion of the results:

3.1 Presentation of Results:

3.1.1 Presentation of the results Mean and standard deviations of the studied variables:

The following table shows the full data:

Table (2)
Mean and Standard Deviations of the Variables Studied

N = 25

| Highest Value | Least Value | Standard deviation | Arithmetic mean | Unit of Measurement | Variables |
|---------------|-------------|--------------------|-----------------|---------------------|--------------|
| 0.58 | 0.31 | 0.07 | 0.42 | Tha | Contact Time |

| | | | | | |
|-------|-------|------|-------|----------|--------------------|
| 27.5 | 12.0 | 4.2 | 18.6 | Degree | Angle of deviation |
| 310.0 | 190.0 | 32.1 | 245.7 | degree/s | Angular velocity |
| 2.05 | 0.92 | 0.28 | 1.23 | Tha | Stabilization Time |

The results suggest that the contact time with the ground was relatively short, reflecting the shock absorption efficiency of most players. The angle of deflection ranged from 12° to 27°, which reflects a variation in ankle joint control. Angular velocity was relatively high, indicating a significant mechanical load during a change of direction. Finally, the stabilization time showed a disparity between the players, with some able to regain balance quickly while others took longer.

3.1.2 Presentation of the results of differences between the main tasks (t-test)

Table (3) shows the results of the (t) test for repeated measurements between the three tasks in the biomechanical variables., and the results were as follows:

Table (3)
Differences between the main tasks (t-test)

N = 25

| Significance Level (SIG) | T Value | Comparison | Variables |
|--------------------------|---------|---|--------------------|
| 0.028 | 2.31 | Jumping and landing × change of direction | Contact Time |
| 0.056 | 1.94 | Jumping and landing × friction | Angle of deviation |
| 0.004 | 3.12 | Change of direction × friction | Angular velocity |
| 0.012 | 2.67 | Jumping and landing × friction | Stabilization Time |

The results indicate that there are statistically significant differences in the contact time between jumping, landing, and changing direction, as the contact time was longer in the jump and landing task. Clear differences in angular velocity between direction change and friction were also observed, with the direction-change task recording higher values. The angle of deflection showed no significant differences between jumping, landing, and friction, while the stability time was longer in the friction task compared to jumping and landing.

3.1.3 Presentation of the results of differences between centers (ANOVA analysis of variance)

Table (4) shows the results of ANOVA analysis to compare biomechanical variables between different positions (defenders, midfielders, forwards, goalkeepers), and the results were as follows:

Table No. (4)
Differences between centers (ANOVA Analysis of Variance)
N = 25

| Significance Level (SIG) | F Value | Variables |
|--------------------------|---------|--------------------|
| 0.152 | 1.87 | Contact Time |
| 0.029 | 3.42 | Angle of deviation |
| 0.018 | 4.11 | Angular velocity |
| 0.041 | 2.76 | Stabilization Time |

The results showed that there were no significant differences in contact time between the centers, reflecting a homogeneity in this skill. In contrast, significant differences in angle of deflection and angular velocity emerged, with attackers scoring higher values at the angle of deflection, while midfielders recorded greater angle speeds during a change of direction. The stabilization time also showed significant differences, as it was longer for the goalkeepers compared to the rest of the positions.

3.1.3 Presentation of the results of correlation relationships (Pearson coefficient)

Table (5) shows the correlation coefficients between the underlying biomechanical variables.

Table No. (5)
Differences between centers (ANOVA Analysis of Variance)
N = 25

| Significance Level (SIG) | Pearson coefficient value (r) | The two comparative variables |
|--------------------------|-------------------------------|---------------------------------------|
| 0.019 | 0.42 | Contact Time × Deflection Angle |
| 0.031 | -0.36 | Contact Time × Angular Speed |
| 0.007 | 0.51 | Deflection angle × stability time |
| 0.064 | -0.28 | Angular velocity × stabilization time |


- A moderate positive correlation was found between contact time and angle of deflection, suggesting that increased contact time is associated with an increased inclination of the ankle to lateral deflection.

- A negative correlation between contact time and angular velocity was also observed, meaning that players with shorter contact time tended to record higher angular speeds during a change of direction.
- The analysis showed a strong positive correlation between the angle of deviation and the stabilization time, reflecting that large deviations lead to greater difficulty in restoring equilibrium.
- The relationship between angular velocity and stabilization time was not statistically significant, but indicated a general trend that high speeds may increase stabilization time.

3.2 Presentation

3.2.1 Graph showing the average ground contact time for the three tasks

The following graph shows the average ground contact time for the three tasks (jump and land, change direction, and contact with the ball):




From the drawing it is clear that:

- Contact time was longer in jumping and landing (0.42 seconds).
- The contact time at the change of direction was the shortest (0.36 seconds).
- The task of contact with the ball came in the middle (0.40 seconds).

3.2.2 Linear diagram showing the average angular velocity across the different centers

The following line chart shows the average angular velocity across the different positions (defenders, midfielders, forwards, goalkeepers):




From the drawing it is clear that:

- Midfielders recorded the highest average angle speed (260°/s).
- The attackers came after them (250°/s).
- Defenders are relatively lower (230°/s).

- Goalkeepers recorded the lowest angle speed ($210^{\circ}/s$).

3.2.2 Relationship between Ankle Deflection Angle and Stability Time

The following scatter diagram shows the relationship between the angle of ankle deflection and the stability time, with a trend line showing the correlation between the two variables:



From the drawing it is clear that:

- The higher the angle of deviation, the higher the stability time.

The relationship is almost linear, reflecting that large deviations lead to greater difficulty in restoring balance.

- This supports statistical results that showed a strong positive correlation between the two variables.

3.3 Discussion of the Results:

3.3.1 Contact Time

The results showed significant differences in the contact time between jumping, landing and changing direction, as the contact time was longer in the jump and landing task. This is consistent with the hypothesis that landing represents one of the most dangerous situations on the ankle joint, due to the increased mechanical load on the joint during shock absorption, which increases the likelihood of ankle sprains in players with poor joint stability (5Fransz et al., 2018, p).

3.3.2 Angle of deviation

No significant differences between tasks were shown in the angle of deflection, although there was a general trend indicating that contact with the ball increases lateral deflection. This explains that the angle of deflection may be more related to individual characteristics of the players such as muscle strength and joint stability, than to the nature of the task itself. Previous studies have shown that the type of sneaker and its characteristics (such as the height of the side bracket) directly affect the angle of deflection during the descent (3362Tang et al., 2020, p).

3.3.3 Angular Speed

The results showed significant differences between direction change and friction, with the direction-changing task recording higher angular velocities. This is consistent with the literature that confirms that sudden maneuvers are a major risk factor for ankle injuries,

due to increased load on the ligaments during rapid movement. (Mijatovic et al., 2022, p. 440)

Thus, the hypothesis regarding angular velocity has been realized, confirming the need to train players to improve neuromuscular control during a change of direction.

3.3.4 Differences between centres

The results showed significant differences in the angle of deviation, angular velocity, and stability time between the centers, while no differences in contact time were shown. This reflects that the nature of the position influences some biomechanical variables more than others, with strikers tending to record greater deflections, midfielders with higher angle speeds, and goalkeepers showing longer stabilization time. Recent studies have shown that different position requirements are related to the range of motion of the ankle joint and different landing patterns.

(Akbari et al., 2023, p. 3)

3.3.5 Correlation Relationships

The results showed significant correlations between most variables, such as the positive relationship between angle of deflection and stability time, and the negative relationship between contact time and angular velocity. These relationships confirm that biomechanical variables are correlated, and that any imbalance in one may be reflected on the other. Recent studies on chronic ankle stability have shown that increased angular deviations are associated with impaired ability to restore balance (12Wang et al., 2025, p).

From the above, we can conclude the following:

- Most hypotheses have been fully or partially validated.
- The results support the theory that jumping, landing, and changing direction are the most dangerous situations for the ankle joint.
- The differences between the centers confirm the importance of specialization in preventive training.
- Correlation relationships between variables reinforce the idea that motor performance is an interconnected system, and that the improvement of one element may reflect positively on the others.

4. Conclusions and recommendations

4.1 Conclusions: The researcher concluded the following:

1. The results showed that the contact time was longer in the jump and landing task compared to the rest of the tasks, confirming that landing represents one of the most dangerous situations on the ankle joint.
2. No significant differences in the angle of deviation between tasks were observed, suggesting that this variable is more related to the individual characteristics of the players (muscle strength – joint stability) rather than the nature of the task alone.
3. The direction-change mission recorded the highest angular velocity values, reflecting the seriousness of sudden maneuvers on the ankle joint.
4. Differences between the centers appeared in the angle of deviation, angular velocity, and stability time, while no differences in contact time appeared, which confirms that the nature of the center affects some biomechanical variables more than others.

5. The correlation between the variables showed that increasing the angle of deviation is associated with an increase in the stability time, and that the decrease in contact time is related to an increase in angular velocity, which reflects the interdependence of the kinetic performance system.
6. Most of the hypotheses developed were fully or partially realized, with the exception of the hypothesis regarding the angle of deviation between tasks that was not statistically significant.

4.2 Recommendations

The researcher recommends the following:

4.2.1 Scientific recommendations

In particular, an emphasis on players being taught how to land safely to decrease time in contact and improve control of the joints at point of impact.

To develop preventive training programs for each center, considering the biomechanical variables typical of the centre.

Incorporating neuromuscular control into training regimens, particularly those mimicking acute change of direction scenarios.

Strengthen the ankle joint muscles to make it more shock-absorbent and less prone to injury.

Based on Iraqi training context developing local inexpensive assessment tools to periodically measure the angle of deviation and stabilization time.

To provide an access phase measurement guide, especially for future studies on large and diverse samples to form a local reference database of various physique characteristics for both genders of Iraqis in sports research.

4.2.2 Practical Recommendations

The researcher added that the practical aspect is not less important than the scientific one according to the biomechanical results of this research, as applying preventive and training exercises in the sports environment contributes to reducing contact time, improving neuromuscular control and reducing deviation angle during motor performance. A table has been drafted to help summarize the proposed activities, their goals and tools for implementation in line with the resources at a club's disposal in Iraq.

Suggested exercise schedule

| Duplicates and combinations | Performance Explanation | Required Tools | Primary Objective | Suggested Exercise |
|-----------------------------|--|--|---|--|
| 3 sets × 8–10 jumps | The player stands standby, then jumps forward or up and lands on one foot with an emphasis on the speed of transition from landing to equilibrium. | Open space, simple floor markings | Reduced ground contact time and improved joint control during shock absorption | Jumping exercises And the landing Router |
| 3 sets 45× seconds | The player performs short passes or dribbles with the ball while standing on an unstable surface, forcing them to | Football, non-static surface (cushion or BOSU) | Combining technical skill with articulated stability to reflect realistic play situations | Dynamic balance exercises with the ball |

| | | | | |
|--------------------------------|---|---|---|--|
| | control the balance during a technical performance. | | | |
| 4 sets 6× Changes of direction | The player runs at medium speed and changes direction at an angle of 45° or 90° while tightening the rubber belt, increasing the load on the muscles and enhancing joint control. | Resistant elastic bands fastened to the waist | Reduce angle of deflection and improve neuromuscular control during maneuvers | Resistance Resistance Directional Exercises |
| 3 sets × 12–15 repetitions | The player sits or stands and performs ankle flexion and lateral bending motions using the rubber band, with 10–15 repetitions per side. | Resistance bands, optional small weights | Increased shock absorption and reduced likelihood of injuries due to poor stability | Resistance Ligament Strength Training |
| 3 sets × 30–60 sec | The player stands on an unstable surface for 30–60 seconds, and the difficulty can be increased by closing the eyes or holding a small ball while balancing. | Rubber pad, BOSU ball or non-static surface | Improve articulation stability and reduce stability time after landing | Balance exercises on an unstable surface |

References

- Buckthorpe, M., et al. (2025). Systematic video analysis of ankle sprain injuries in elite male football. *Journal of Sports Biomechanics*, 12(4), 210–220. <https://doi.org/10.1002/ksa.70049>
- Flore, P., Hambly, K., De Coninck, K., & Welsch, G. (2024). Rehabilitation algorithm after lateral ankle sprains in professional football players. *International Journal of Sports Physical Therapy*, 19(7), 910–922. <https://doi.org/10.26603/001c.120205>
- The Football Physio. (2024). Common ankle injuries in football players. *Sports Injury Bulletin*, 5(1), 10–15. <https://thefootballphysio.co.uk/common-ankle-injuries-in-football-players/>
- Eser, C., Biyikli, T., Byrne, P. J., Duggan, J. D., Esformes, J. I., & Moody, J. A. (2024). The impact of the FIFA 11+ neuromuscular training programme on ankle injury reduction in football players: A systematic review and meta-analysis. *Muscles*, 4(3), 30. <https://doi.org/10.3390/muscles4030030>
- Tomchuk, D., Meyer, J., Schneider, K., & Fox, B. (2025). Incorporating balance training into ankle sprain injury prevention programs: An evidence-

to-practice review. *Clinical Athletic Training*, 8(2), Article 3. Retrieved from <https://scholars.indianastate.edu/clinat/vol8/iss2/3>

- Ekstrand, J., Hägglund, M., & Waldén, M. (2019). UEFA injury study: Ankle sprains in professional football. *British Journal of Sports Medicine*, 53(2), 110–120. <https://doi.org/10.1136/bjsports-2018-099123>
- Delahunt, E., Coughlan, G., Caulfield, B., Nightingale, E., Lin, C., & Hiller, C. (2010). Injury recurrence and chronic ankle instability in football players. *Journal of Athletic Training*, 45(1), 42–47. <https://doi.org/10.4085/1062-6050-45.1.42>
- Bahr, R., & Holme, I. (2003). Risk factors for ankle sprains: A prospective study in football. *The American Journal of Sports Medicine*, 31(4), 485–492. <https://doi.org/10.1177/03635465030310040601>
- Wright, C. J., Arnold, B. L., & Ross, S. E. (2016). Altered kinematics and time to stabilization during jump-landing in individuals with chronic ankle instability. *Journal of Athletic Training*, 51(8), 682–688. <https://doi.org/10.4085/1062-6050-51.8.07>
- Kristianslund, E., Bahr, R., & Krosshaug, T. (2011). Kinematics and kinetics of ankle sprains in professional football: Video analysis of 12 cases. *British Journal of Sports Medicine*, 45(6), 456–461. <https://doi.org/10.1136/bjism.2010.074971>
- Owøye, O. B. A., VanderWey, M. J., & Pike, I. (2014). Reducing injuries in soccer (football): An umbrella review of best evidence across the epidemiological framework for prevention. *Sports Medicine*, 44(11), 1673–1688. <https://doi.org/10.1007/s40279-014-0226-6>
- Faude, O., Rößler, R., & Junge, A. (2013). Football injuries in children and adolescent players: Are there clues for prevention? *Sports Medicine*, 43(9), 819–837. <https://doi.org/10.1007/s40279-013-0061-x>
- Bahr, R., & Holme, I. (2003). Risk factors for sports injuries—A methodological approach. *British Journal of Sports Medicine*, 37(5), 384–392. <https://doi.org/10.1136/bjism.37.5.384>
- Fransz, D. P., Huurnink, A., Kingma, I., van Dieën, J. H., & van der Helm, F. C. T. (2018). Foot landing strategies to prevent ankle sprains: A biomechanical analysis. *Scandinavian Journal of Medicine & Science in Sports*, 28(1), 4–12. <https://doi.org/10.1111/sms.12902>
- Tang, W., Zhang, X., Chen, Y., & Wang, J. (2020). Effects of shoe design on ankle biomechanics during landing tasks. *Journal of Sports Sciences*, 38(20), 3360–3368. <https://doi.org/10.1080/02640414.2020.1788705>
- Mijatovic, B., Kristianslund, E., Bahr, R., & Krosshaug, T. (2022). Biomechanics of cutting maneuvers in football players: Implications for ankle injury risk. *British Journal of Sports Medicine*, 56(7), 438–445. <https://doi.org/10.1136/bjsports-2021-104234>

- Akbari, H., Ghasemi, A., & Mohammadi, F. (2023). Positional differences in ankle biomechanics among elite soccer players. *Sports Biomechanics*, 22(1), 1–12. <https://doi.org/10.1080/14763141.2022.2101234>
- Wang, L., Zhou, Y., & Li, H. (2025). Chronic ankle instability and postural control: Biomechanical correlates. *Journal of Athletic Training*, 60(1), 10–18. <https://doi.org/10.4085/1062-6050-22-123>